

On Monotonocities of the Generalized Mean Ratio and Related Results

LEON JAY GLESER*

Johns Hopkins University, Baltimore, Maryland

Submitted by R. Bellman

1. INTRODUCTION AND SUMMARY

The purpose of this paper is to discuss lower (and upper) bounds for ratios of weighted means M_r , $-\infty \leq r \leq \infty$, of positive variables x_1, x_2, \dots, x_n in terms of corresponding ratios of weighted means for any subset $\{x_{j_1}, x_{j_2}, \dots, x_{j_m}\}$, $m < n$, of these values. Some of the results obtained are extended to a more abstract context.

2. A MONOTONICITY OF THE GENERALIZED MEAN RATIO

Let p_1, p_2, \dots, p_n ($n > 1$) be positive numbers with $\sum_{i=1}^n p_i = 1$. For every collection x_1, x_2, \dots, x_n of positive real numbers, and for every real v , define the v -th weighted mean M_v to be

$$M_v = \left(\sum_{i=1}^n p_i x_i^v \right)^{1/v}, \quad (2.1)$$

where it is understood that

$$\lim_{v \rightarrow \infty} M_v = \max_{1 \leq i \leq n} x_i, \quad \lim_{v \rightarrow -\infty} M_v = \min_{1 \leq i \leq n} x_i,$$

and

$$\lim_{v \rightarrow 0} M_v = \prod_{i=1}^n x_i^{p_i}.$$

* Research sponsored by the Air Force Office of Scientific Research, AFOSR Contract No. AF 49(638)-1302.

The *generalized mean ratio* $T_{r,s}$ is defined to be

$$T_{r,s}(n) = \frac{M_r}{M_s}, \quad r > s. \quad (2.2)$$

The well known *moment inequality* (viz., [1]) tells us that $T_{r,s}(n) \geq 1$, with strict inequality unless $x_1 = x_2 = \cdots = x_n$.

Suppose that we eliminate one of the x_i 's (say, x_{j_0}) and renormalize the probabilities p_i , $i \neq j_0$, so that they sum to one. Form the reduced generalized mean ratio

$$T_{r,s}(n-1; j_0) \equiv \left[\frac{\sum_{i \neq j_0} p_i x_i^r}{\sum_{i \neq j_0} p_i} \right]^{1/r} \left[\frac{\sum_{i \neq j_0} p_i x_i^s}{\sum_{i \neq j_0} p_i} \right]^{-1/s} \quad (2.3)$$

from the remaining x_i 's, $i \neq j_0$. It might be hypothesized that $T_{r,s}(n) \geq T_{r,s}(n-1; j_0)$ for every j_0 (i.e., that the generalized mean ratio is monotone in n). This hypothesis is false in general; however, we can verify the weaker assertion: $T_{r,s}(n) \geq T_{r,s}(n-1; j)$, for some j .

THEOREM 2.1. If $r > s$,

$$T_{r,s}(n) \geq \min_{1 \leq j \leq n} T_{r,s}(n-1; j). \quad (2.4)$$

PROOF. Let

$$q_j = \sum_{i \neq j} p_i, \quad w_j^s = q_j^{-1} \sum_{i \neq j} p_i x_i^s.$$

Then

$$\begin{aligned} T_{r,s}(n) &= \frac{\left[(n-1)^{-1} \sum_{j=1}^n q_j w_j^r (T_{r,s}(n-1; j))^r \right]^{1/r}}{\left[(n-1)^{-1} \sum_{j=1}^n q_j w_j^s \right]^{1/s}} \\ &\geq \left[\min_{1 \leq j \leq n} T_{r,s}(n-1; j) \right] \frac{\left\{ (n-1)^{-1} \sum_{j=1}^n q_j w_j^r \right\}^{1/r}}{\left\{ (n-1)^{-1} \sum_{j=1}^n q_j w_j^s \right\}^{1/s}}. \end{aligned} \quad (2.5)$$

But $(n-1)^{-1} \sum_{j=1}^n q_j = 1$, so that the result (2.4) follows from the moment inequality

$$\left\{ (n-1)^{-1} \sum_{j=1}^n q_j w_j^r \right\}^{1/r} \geq \left\{ (n-1)^{-1} \sum_{j=1}^n q_j w_j^s \right\}^{1/s}. \quad ||$$

We might also ask whether there is always a j_0 such that $T_{r,s}(n-1; j_0) \geq T_{r,s}(n)$. In general, this assertion is false. The following easily verified condition, however, is sufficient for its validity.

THEOREM 2.2. *Let $x_{[1]} \leq x_{[2]} \leq \dots \leq x_{[n]}$ be the ordered values of the x_i 's. If $r > s$, and if for some k , $1 \leq k \leq n$, we have*

$$M_s \leq x_{[k]} \leq M_r, \quad (2.6)$$

then

$$T_{r,s}(n) \leq \max_{1 \leq j \leq n} T_{r,s}(n-1; j).$$

Proof. Let $p_{[i]}$ be the probability associated with $x_{[i]}$, $i = 1, 2, \dots, n$. From (2.6),

$$M_r^r \leq \left(\sum_{i \neq k} p_{[i]} x_{[i]}^r \right) + p_{[k]} M_r^r,$$

$$M_s^s \geq \left(\sum_{i \neq k} p_{[i]} x_{[i]}^s \right) + p_{[k]} M_s^s.$$

Thus

$$M_r \leq \left((1 - p_{[k]})^{-1} \sum_{i \neq k} p_{[i]} x_{[i]}^r \right)^{1/r} \equiv M_r([k]),$$

$$M_s \geq \left((1 - p_{[k]})^{-1} \sum_{i \neq k} p_{[i]} x_{[i]}^s \right)^{1/s} \equiv M_s([k]),$$

and consequently

$$T_{r,s} = \frac{M_r}{M_s} \leq \frac{M_r([k])}{M_s([k])} \leq \max_{1 \leq j \leq n} T_{r,s}(n-1; j). \quad ||$$

Similar results can be obtained for comparing the generalized mean ratio $T_{r,s}(n)$ with reduced generalized mean ratios formed from arbitrary subcollections $\{x_{j_1}, x_{j_2}, \dots, x_{j_m}\}$, $m < n$, of $\{x_1, x_2, \dots, x_n\}$. Such assertions, and

many other statements of interest to us, can be very conveniently obtained as special cases of results found in the generalization of our problem to a more abstract, measure-theoretic context.

3. MONOTONICITIES OF THE GENERALIZED MEAN RATIO ON ARBITRARY PROBABILITY SPACES

Consider the probability space $(\mathcal{X}, \mathcal{B}, P)$, where P is a probability measure defined on the σ -algebra \mathcal{B} of subsets of \mathcal{X} . Consider any positive measurable function f , and define the v -th moment of f to be

$$\mu_v(\mathcal{X}) = \int_{\mathcal{X}} f^v(x) dP(x), \quad -\infty \leq v \leq \infty. \quad (3.1)$$

Then the generalized mean ratio defined with respect to f is

$$T_{r,s}(\mathcal{X}) \equiv \frac{\mu_r^{1/r}}{\mu_s^{1/s}}, \quad r > s. \quad (3.2)$$

We note that $\lim_{r \rightarrow \infty} \mu_r^{1/r}$ can be taken to be the essential supremum of $f(x)$ over \mathcal{X} and that $\lim_{s \rightarrow -\infty} \mu_s^{1/s}$ can be taken to be the essential infimum of f over \mathcal{X} . Further, $\lim_{r \rightarrow 0} \mu_r^{1/r}$ equals $\exp\{\int_{\mathcal{X}} \log f(x) dP(x)\}$. Note also that the moment inequality in this context states that $T_{r,s}(\mathcal{X}) \geq 1$, with equality holding if and only if f is essentially (almost surely) constant.

Consider, now, any subset $A \in \mathcal{B}$ of \mathcal{X} . Renormalize the probability measure P over A so that the resulting measure is again a probability measure. Such normalization yields the conditional probability measure:

$$P(B | A) \equiv \begin{cases} P(B \cap A)/P(A), & \text{if } P(A) > 0, \\ 0, & \text{if } P(A) = 0, \end{cases} \quad (3.3)$$

all $B \in \mathcal{B}$. Define the v th conditional moment of f to be

$$\mu_v(A) = \int_A f^v(x) dP(x | A), \quad (3.4)$$

and define the generalized mean ratio of f with respect to $P(\cdot | A)$ to be

$$T_{r,s}(A) = \frac{\mu_r^{1/r}(A)}{\mu_s^{1/s}(A)}, \quad r > s. \quad (3.5)$$

The discrete case discussed in Section 2 is a special case of the above model. For, if $\mathcal{X} = \{x_1, x_2, \dots, x_n\}$, P is the measure putting mass p_i on i ,

$i = 1, 2, \dots, n$, \mathcal{B} is the collection of all subsets $A = \{j_1, j_2, \dots, j_m\}$ of \mathcal{X} , and $f(i) = x_i$, $i = 1, 2, \dots, n$, then $T_{r,s}(\mathcal{X})$ corresponds to $T_{r,s}(n)$. Further, if $A_j = \{i : i \neq j\}$, then $T_{r,s}(A_j) = T_{r,s}(n-1; j)$.

We have already seen in Section 2 that $T_{r,s}(\mathcal{X})$ need not exceed $T_{r,s}(A)$ for $A \in \mathcal{B}$. The assertion that $T_{r,s}(\mathcal{X}) \geq \min_{A \in \mathcal{B}} T_{r,s}(A)$ is meaningless if we allow \mathcal{B} to contain point sets $A_\epsilon = \{x : f(x) = \epsilon\}$. For example, if for some ϵ_0 , $P(A_{\epsilon_0}) > 0$, the assertion $T_{r,s}(\mathcal{X}) \geq \min_{A \in \mathcal{B}} T_{r,s}(A)$ becomes $T_{r,s}(\mathcal{X}) \geq T_{r,s}(A_{\epsilon_0}) = 1$, which tells us no more than the moment inequality. We can, however, ask for sufficient conditions for $T_{r,s}(\mathcal{X}) \geq T_{r,s}(A)$, given $A \in \mathcal{B}$.

LEMMA 3.1. *If $r > s$, $A \in \mathcal{B}$, then*

$$\min \{T_{r,s}(A), T_{r,s}(A^c)\} \leq T_{r,s}(\mathcal{X}) \leq [\max \{T_{r,s}(A), T_{r,s}(A^c)\}] \frac{[p + (1-p)\gamma^r]^{1/r}}{[p + (1-p)\gamma^s]^{1/s}}, \quad (3.6)$$

where A^c is the complement of A in \mathcal{X} , $p = P(A) = 1 - P(A^c)$, and $\gamma^s = \mu_s(A^c)/\mu_s(A)$.

PROOF. Note that

$$\begin{aligned} T_{r,s}(\mathcal{X}) &= \frac{[p\mu_r(A) + (1-p)\mu_r(A^c)]^{1/r}}{[p\mu_s(A) + (1-p)\mu_s(A^c)]^{1/s}} \\ &= \frac{[p(T_{r,s}(A))^r + (1-p)\gamma^r(T_{r,s}(A^c))^r]^{1/r}}{[p + (1-p)\gamma^s]^{1/s}}. \end{aligned} \quad (3.7)$$

Since the moment inequality tells us that

$$t(\gamma) \equiv \frac{(p + (1-p)\gamma^r)^{1/r}}{(p + (1-p)\gamma^s)^{1/s}} \geq 1,$$

and since

$$t(\gamma) \min \{T_{r,s}(A), T_{r,s}(A^c)\} \leq T_{r,s}(\mathcal{X}) \leq t(\gamma) \max \{T_{r,s}(A), T_{r,s}(A^c)\},$$

the result (3.6) follows. \parallel

THEOREM 3.2. *If $r > s$, $A \in \mathcal{B}$, then*

$$T_{r,s}(A^c) \geq T_{r,s}(A) \Rightarrow T_{r,s}(\mathcal{X}) \geq T_{r,s}(A), \quad (3.8)$$

$$T_{r,s}(A^c) \geq \Delta^* T_{r,s}(A) \Leftrightarrow T_{r,s}(\mathcal{X}) \geq T_{r,s}(A), \quad (3.9)$$

where Δ^* is the solution of

$$(p + (1-p)\gamma^r(\Delta^*)^r)^{1/r} = (p + (1-p)\gamma^s)^{1/s}. \quad (3.10)$$

PROOF. The result (3.8) is a direct consequence of Lemma 3.1. To prove (3.9), note that from (3.7) we have

$$T_{r,s}(\mathcal{X}) = T_{r,s}(A) \frac{[p + (1-p)\gamma^r \Delta^r]^{1/r}}{[p + (1-p)\gamma^s]^{1/s}},$$

where $\Delta = T_{r,s}(A^c)/T_{r,s}(A)$. Thus $T_{r,s}(\mathcal{X})$ exceeds $T_{r,s}(A)$ if and only if

$$\frac{[p + (1-p)\gamma^r \Delta^r]^{1/r}}{[p + (1-p)\gamma^s]^{1/s}} \geq 1. \quad (3.11)$$

It is easy to see that (3.11) holds if and only if $\Delta \geq \Delta_y^*$, where Δ_y^* is the solution of (3.10). \parallel

4. A GENERAL BOUND

In Section 3, our investigations of the relations between $T_{r,s}(\mathcal{X})$ and $T_{r,s}(A)$ require us to know $T_{r,s}(A^c)$ and $\gamma^s = \mu_s(A^c)/\mu_s(A)$. Since $T_{r,s}(A^c)$ and γ^s depend not only on A and $P(\cdot)$ for their values, but also on the nature of f , the results of Section 3 may be difficult to apply.

If instead of asking for a bound on $T_{r,s}(\mathcal{X})$ of the form $T_{r,s}(\mathcal{X}) \geq T_{r,s}(A)$, we ask for a bound of the form

$$T_{r,s}(\mathcal{X}) \geq h_p(T_{r,s}(A)) \geq 1,$$

where $h_p(u)$ is a function defined on $[1, \infty)$ with form depending on A only through $p \equiv P(A)$, the following lemma provides one solution to our problem.

LEMMA 4.1. For $r > s$,

$$T_{r,s}(\mathcal{X}) \geq \frac{T_{r,s}(A)}{[p + (1-p)(T_{r,s}(A))^t]^{1/t}} \geq 1, \quad (4.1)$$

where $t = rs/(r-s)$.

PROOF. From Eq. (3.7) and the fact that $T_{r,s}(A^c) \geq 1$, we have

$$T_{r,s}(\mathcal{X}) \geq \frac{[p\mu_r(A) + (1-p)x^r]^{1/r}}{[p\mu_s(A) + (1-p)x^s]^{1/s}}, \quad (4.2)$$

where $x^s = \mu_s(A^c)$. It is easy to verify (4.1) for $p = P(A) = 0, 1$. Consequently, we can assume $0 < p < 1$ in what follows. Differentiating the right-hand expression with respect to x , we find that the sign of the derivative is equal to the sign of $x^{r-s} - (\mu_r(A))^r (\mu_s(A))^{-s}$. Thus $T_{r,s}(\mathcal{X})$ is minimized only at $(x^*)^{r-s} = (\mu_r(A))^r (\mu_s(A))^{-s}$. Substituting x^* in (4.2) gives us (4.1). \parallel

COROLLARY 4.2. For $r > s$, any j , $1 \leq j \leq n$,

$$T_{r,s}(n) \geq \frac{T_{r,s}(n-1; j)}{(q_j + p_j(T_{r,s}(n-1; j))^t)^{1/t}}, \quad (4.3)$$

where

$$t = \frac{rs}{r-s}, \quad q_j = \sum_{i \neq j} p_i.$$

If $r > 0$, $s \leq 0$, we can obtain a somewhat more appealing lower bound for $T_{r,s}(\mathcal{A})$ in terms of $T_{r,s}(A)$ and $p = P(A)$.

LEMMA 4.3. For $r > 0$, $s \leq 0$, $A \in \mathcal{B}$,

$$T_{r,s}(\mathcal{A}) \geq T_{r,s}^p(A) \geq 1. \quad (4.4)$$

PROOF. If $r > 0$, $s \leq 0$, then $t = rs/r - s \leq 0$. Consequently, the moment inequality tells us that

$$[p + (1-p)(T_{r,s}(A))^t]^{1/t} \leq [T_{r,s}(A)]^{(1-p)}. \quad (4.5)$$

Equation (4.4) now follows from (4.3) and (4.4). \parallel

COROLLARY 4.4. For $r > 0$, $s \leq 0$, any j , $1 \leq j \leq n$,

$$T_{r,s}(n) \geq (T_{r,s}(n-1; j))^{q_j}, \quad (4.6)$$

where

$$q_j = \sum_{i \neq j} p_i.$$

REMARK. If $r > s$, $rs > 0$, the results (4.5) and (4.6) do not, in general, hold.

ACKNOWLEDGMENT

I am grateful to A. W. Marshall and I. Olkin for their comments and suggestions.

REFERENCES

1. G. A. HARDY, J. E. LITTLEWOOD AND G. POLYA. "Inequalities," 2nd ed., Cambridge University Press, Cambridge, 1959.